

A DICHOTOMY FOR BOREL FUNCTIONS

MARCIN SABOK

ABSTRACT. The dichotomy discovered by Solecki in [3] states that any Baire class 1 function is either σ -continuous or “includes” the Pawlikowski function P . The aim of this paper is to give an argument which is simpler than the original proof of Solecki and gives a stronger statement: a dichotomy for all Borel functions.

1. INTRODUCTION

An old question of Lusin asked whether there exists a Borel function which cannot be decomposed into countably many continuous functions. By now several examples have been given, by Keldiš, Adyan and Novikov among others. A particularly simple example, the function $P : (\omega + 1)^\omega \rightarrow \omega^\omega$, has been found by Pawlikowski (cf. [1]). By definition,

$$P(x)(n) = \begin{cases} x(n) + 1 & \text{if } x(n) < \omega, \\ 0 & \text{if } x(n) = \omega. \end{cases}$$

It is proved in [1] that if $A \subseteq (\omega + 1)^\omega$ is such that $P \upharpoonright A$ is continuous then $P[A] \subseteq \omega^\omega$ is nowhere dense. Since P is a surjection, it is not σ -continuous.

In [3] Solecki showed that the above function is, in a sense, the only such example, at least among Baire class 1 functions (in other words, it is the initial object in a certain category).

Theorem 1 (Solecki, [3]). *For any Baire class 1 function $f : X \rightarrow Y$, where X, Y are Polish spaces, either f is σ -continuous or there exist topological embeddings φ and ψ such that the following diagram commutes:*

$$\begin{array}{ccc} \omega^\omega & \xrightarrow{\psi} & Y \\ \uparrow P & & \uparrow f \\ (\omega + 1)^\omega & \xrightarrow{\varphi} & X \end{array}$$

In [4] Zapletal generalized Solecki’s dichotomy to all Borel functions by proving the following theorem.

Theorem 2 (Zapletal, [4]). *If $f : X \rightarrow Y$ is a Borel function which is not σ -continuous then there is a compact set $C \subseteq X$ such that $f|_C$ is not σ -continuous and of Baire class 1.*

In this paper we give a new proof of the above dichotomy for all Borel functions, which is direct, shorter and more general than the original proof from [3].

2. NOTATION

We say that a Borel function $f : X \rightarrow Y$, where X, Y are Polish spaces, is σ -continuous if there exist a countable cover of the space $X = \bigcup_n X_n$ (with arbitrary sets X_n) such that $f|_{X_n}$ is continuous for each n . It follows from the Kuratowski extension theorem that we may require that the sets X_n be Borel. If f is a Borel function which is not σ -continuous then the family of sets on which it is σ -continuous is a proper σ -ideal in X . We denote this σ -ideal by I_f .

In a metric space (X, d) for $A, B \subseteq X$ let us denote by $h(A, B)$ the Hausdorff distance between A and B .

The spaces $(\omega + 1)^\omega$ and $(\omega + 1)^n$ for $n < \omega$ are endowed with the product topology of order topologies on $\omega + 1$.

3. THE ZAPLETAL'S GAME

In [4] Zapletal introduced a two-player game, which turns out to be very useful in examining σ -continuity of Borel functions. Let $B \subseteq \omega^\omega$ be a Borel set and $f : B \rightarrow 2^\omega$ be a Borel function. Let $\rho : \omega \rightarrow \omega \times 2^{<\omega} \times \omega$ be a bijection. The game $G_f(B)$ is played by Adam and Eve. They take turns playing natural numbers. In his n -th move, Adam picks $x_n \in \omega$. In her n -th move, Eve chooses $y_n \in 2$. At the end of the game we have $x \in \omega^\omega$ and $y \in 2^\omega$ formed by the numbers picked by Adam and Eve, respectively. Next, $y \in 2^\omega$ is used to define a sequence of partial continuous functions (with domains of type G_δ in ω^ω) in the following way. For $n < \omega$ let f_n be a partial function from ω^ω to 2^ω such that for $t \in \omega^\omega$ and $\sigma \in 2^{<\omega}$

$$f_n(t) \supseteq \sigma \quad \text{iff} \quad \exists k \in \omega \quad y(\rho(n, \sigma, k)) = 1$$

and $\text{dom}(f_n) = \{t \in \omega^\omega : \forall n < \omega \exists! \sigma \in 2^n \ f_n(t) \supseteq \sigma\}$. Eve wins the game $G_f(B)$ if $x \notin B$ or $\exists n \ f(x) = f_n(x)$. Otherwise Adam wins the game.

It is easy to see that if f is a Borel function then G_f is a Borel game. The key feature of the game G_f is that it detects σ -continuity of the function f .

Theorem 3 (Zapletal,[4]). *For $B \subseteq \omega^\omega$ and $f : B \rightarrow 2^\omega$ Eve has a winning strategy in the game $G_f(B)$ if and only if f is σ -continuous on B .*

Note that if Adam has a winning strategy then the image of his strategy (treated as a continuous function from 2^ω to B) is a compact set on which f is also not σ -continuous. This observation and the Borel determinacy gives the following corollary.

Corollary 1 (Zapletal,[4]). *If B is a Borel set and $f : B \rightarrow 2^\omega$ is a Borel function which is not σ -continuous then there is a compact set $C \subseteq B$ such that $f|_C$ is also not σ -continuous.*

4. PROOF OF THE DICHOTOMY

In the statement of Theorem 1 both functions φ and ψ are to be topological embeddings. However, as we will see below, for the dichotomy it is enough that they both are injective, φ continuous and ψ open. We are going to prove first this version of the dichotomy.

Theorem 4. *Let X be a Polish space and $f : X \rightarrow 2^\omega$ be a Borel function. Then precisely one of the following conditions holds:*

- (1) *either f is σ -continuous*
- (2) *or there are an open injection ψ and a continuous injection φ such that the following diagram commutes:*

$$\begin{array}{ccc} \omega^\omega & \xrightarrow{\psi} & 2^\omega \\ \uparrow P & & \uparrow f \\ (\omega + 1)^\omega & \xrightarrow{\varphi} & X \end{array}$$

Notice that compactness of $(\omega + 1)^\omega$ implies that the ψ above must be a topological embedding.

Proof. It is straightforward that (2) implies that f is not σ -continuous. Let us assume that f is not σ -continuous and prove that (2) holds. By Corollary 1 we may assume that X is compact.

Notation. First we introduce some notation. For a fixed n and $0 \leq k \leq n$ let S_k^n be the set of points in $(\omega + 1)^n$ of Cantor-Bendixson rank $\geq n - k$. For each $n < \omega$ and $1 \leq k \leq n$ let us pick a function $\pi_k^n : S_k^n \rightarrow S_{k-1}^n$ such that

- on S_{k-1}^n π_k^n is the identity,

- if $\tau \in S_k^n \setminus S_{k-1}^n$ then we pick one $i \in n$ such that $\tau(i) < \omega$ and $\tau(i)$ is maximal such and define

$$\pi_k^n(\tau)(i) = \omega, \quad \pi_k^n(\tau)(j) = \tau(j) \quad \text{for } j \neq i.$$

This definition clearly depends on the choice of the index i above. Note, however, that we may pick the functions π_k^n so that they are coherent, in the sense that for $\tau \in (\omega+1)^{n+1}$, unless $\tau(n)$ is the biggest finite value of τ , we have $\pi_{k+1}^{n+1}(\tau) = \pi_k^n(\tau \upharpoonright n) \cap \tau(n)$. In particular $\pi_{k+1}^{n+1}(\sigma \cap \omega) = \pi_k^n(\sigma) \cap \omega$ for any $\sigma \in (\omega+1)^n$. The functions π_k^n will be called projections.

Lemma 1. *For each n and $1 \leq k \leq n$ the projection $\pi_k^n : S_k^n \rightarrow S_{k-1}^n$ is continuous.*

Proof. Note that any point in S_k^n except $(\omega, \dots, \omega)$ (k times ω) has a neighborhood in which projection is unambiguous and hence continuous. But it is easy to see that at the point $(\omega, \dots, \omega)$ any projection is continuous. $\square$

For each $n < \omega$ let us also introduce the function $r_n : (\omega+1)^n \rightarrow (\omega+1)^n$ defined as $r_n(\tau \cap a) = \tau \cap \omega$.

To make the above notation more readable we will usually drop subscripts and superscripts in π_k^n and r_n .

We pick a well-ordering $\leq$ of $(\omega+1)^{<\omega}$ into type ω such that for each point $\tau \in (\omega+1)^{<\omega}$ all elements of the transitive closure of τ with respect to π , r and restrictions (i.e. functions of the form $(\omega+1)^n \ni \tau \mapsto \tau \upharpoonright m \in (\omega+1)^m$ for $m < n$) are $\leq \tau$.

For a set $B \in \text{Bor}(X) \setminus I_f$ let B^* denote the set B shrunk by all basic clopens C which have I -small intersection with B .

Strategy of the construction. In order to define functions φ and ψ , we will construct for each $\tau \in (\omega+1)^{<\omega}$ a clopen set $C_\tau \subseteq 2^\omega$ and a compact set $X_\tau \subseteq X$ such that if $\sigma \subseteq \tau$ then $C_\tau \subseteq C_\sigma$ and $X_\tau \subseteq X_\sigma$.

The sets C_τ will be disjoint, which means that for $\tau \neq \tau'$, $|\tau| = |\tau'|$ $C_\tau \cap C_{\tau'} = \emptyset$. We will also need $X_\tau \subseteq f^{-1}[C_\tau]$ and $\text{diam}(X_\tau) < 1/|\tau|$.

The construction of the sets X_τ, C_τ will be done by induction along the ordering $\leq$ on $(\omega+1)^{<\omega}$. In fact, we will do something more: at each step n if τ is the n -th element of $(\omega+1)^{<\omega}$ we will construct not only a compact set X_τ but also I_f -positive Borel sets X_σ^n for $\sigma \leq \tau$ such that:

- $X_\tau \subseteq X_{\tau \upharpoonright (|\tau|-1)}^{n-1}$,
- $X_\sigma^n \subseteq X_\sigma^{n-1}$ if $\sigma < \tau$,
- $X_{\sigma \cap a}^n \subseteq X_\sigma^n$ if $\sigma, \sigma \cap a < \tau$,
- $X_\sigma^n \cap f^{-1}[C_{\sigma \cap a}] = \emptyset$ if $\sigma, \sigma \cap a \leq \tau$,

- $X_{\sigma^\frown \omega}^n \subseteq \text{cl}(X_\sigma^n)$ if $\sigma, \sigma^\frown \omega \leq \tau$.

The set X_σ^n is to be understood as the space for further construction of sets X_ρ for $\rho \supseteq \sigma$ and $\rho > \tau$, as can be seen in the first condition above. The last condition, as we will see later, will be used to guarantee “continuity” of the family of sets X_τ . For technical reasons we will also make sure that $X_\sigma^n = (X_\sigma^n)^*$.

We are going to ensure disjointness of C_τ ’s by satisfying the following conditions:

- $C_{\tau^\frown a} \subseteq C_\tau$,
- $C_{\tau^\frown a} \cap C_{\tau^\frown b} = \emptyset$ for $a \neq b$.

The fact that $\text{diam}(X_\tau) < 1/|\tau|$ will follow from the following inductive conditions (recall that $\pi(\tau) \leq \tau$ for any τ):

- $\text{diam}(X_\tau) < 3 \text{diam}(X_{\pi(\tau)})$,
- $\text{diam}(X_{\tau^\frown \omega}) < 1/(3^{|\tau|+1}(|\tau| + 1))$,

because iterating projections in $(\omega + 1)^n$ stabilizes before $n + 1$ steps.

The crucial feature of the sets X_τ is that this family should be “continuous”. Namely, we will require that if τ and $\pi(\tau)$ occur by the n -th step then

$$(1) \quad h(X_\tau^n, X_{\pi(\tau)}^n) < 3^{|\tau|} d(\tau, \pi(\tau))$$

This condition is the most difficult. To fulfill it we will construct yet another kind of objects. Notice first that if $h(A, B) < \varepsilon$ for two nonempty sets in X then there are two finite families (we will refer to them as to “anchors”) A_i and B_i ($i \in I_0$) of subsets of A and B respectively such that for any $A'_i \subseteq A_i$, $B'_i \subseteq B_i$ still $h(\bigcup_i A'_i, \bigcup_i B'_i) < \varepsilon$. Similarly, if $h(A, B) < \varepsilon$ and $C \subseteq A$ then there exist a finite family D_i ($i \in I_0$) of subsets of B such that for any $D'_i \subseteq D_i$ $h(\bigcup_i D'_i, C) < \varepsilon$.

At each step n if τ is the n -th element of $(\omega + 1)^{<\omega}$ we will additionally construct anchors

- for each pair X_σ^n and $X_{\pi(\sigma)}^n$ such that $\sigma, \pi(\sigma) \leq \tau$
- and for each tripple $X_\sigma^n, X_{\pi(\sigma)}^n, X_{\sigma^\frown a}^n$ such that $a \in \omega + 1$, $\pi(\sigma^\frown a) \subseteq \pi(\sigma)$ and $\sigma, \pi(\sigma), \sigma^\frown a \leq \tau$.

Completing the diagram. As we now have a clear picture of what should be constructed let us argue that this is enough to finish the proof. For each $t \in (\omega + 1)^\omega$ the intersection $\bigcap_n X_{t|n}$ has precisely one point so let us define $\varphi(t)$ to be this point. The other function, ψ is defined as $f \circ \varphi \circ P^{-1}$. Let us check that this works. Both functions ψ and φ are injective thanks to the disjointness of the sets C_τ and to the fact that $X_\tau \subseteq f^{-1}[C_\tau]$. The function ψ is open because C_τ are clopens.

To see continuity of φ notice first that since the sets X_τ have diameters vanishing to 0, it suffices to check that φ is continuous on each $(\omega + 1)^n$ (which are treated as subsets of $(\omega + 1)^\omega$ via the embedding $e : \tau \mapsto \tau^\frown(\omega, \omega, \dots)$). Continuity on $(\omega + 1)^n$ is checked inductively on the sets S_k^n for $0 \leq k \leq n$.

The set S_0^n consists of one point, so there is nothing to check. Suppose that $\tau_i \rightarrow \tau$, $\tau, \tau_i \in S_k^n, i \in \omega$. Then either the sequence is eventually constant or $\tau \in S_{k-1}^n$. Let us assume the latter. By the inductive assumption and continuity of projection $\varphi(\pi(\tau_i)) \rightarrow \varphi(\tau)$. Now pick any $\varepsilon > 0$. Let m be such that $\text{diam}(X_\sigma) < \varepsilon$ for $\sigma \in (\omega + 1)^m$ and $j \in \omega$ such that $d(\tau_j, \pi(\tau_j)) < 3^{-m}\varepsilon$. Let us write $\rho^\frown\omega^l$ for ρ extended by l many ω 's. By (1) and coherence of projections we have

$$h(X_{\tau_j^\frown\omega^{m-n}}, X_{\pi(\tau_j)^\frown\omega^{m-n}}) < \varepsilon,$$

which implies that $\varphi(\tau_j)$ and $\varphi(\pi(\tau_j))$ are closer than 3ε . This shows that $\varphi(\tau_j) \rightarrow \varphi(\tau)$ and proves continuity of φ .

Key lemma. Now we state the key lemma, which will be used to guarantee “continuity” of the family of sets X_τ .

Lemma 2. *Let X be a Borel set, $f : X \rightarrow \omega^\omega$ a Borel, not σ -continuous function. There exist a basic clopen $C_\omega \subseteq f[X]$ and a compact set $X_\omega \subseteq f^{-1}[C_\omega]$ such that*

- $f|_{X_\omega}$ is not σ -continuous,
- $X_\omega \subseteq \text{cl}((f^{-1}[\omega^\omega \setminus C_\omega])^*)$.

The compact set X_ω can be chosen of arbitrarily small diameter.

Proof. Without loss of generality assume that $f^{-1}[C] = (f^{-1}[C])^*$ for all clopen sets $C \subseteq \omega^\omega$. Let us consider the following tree of open sets, indexed by $\omega^{<\omega}$

$$U_\tau = \text{int}(f^{-1}[[\tau]]).$$

Let $G = \bigcap_n \bigcup_{|\tau|=n} U_\tau$ and $Z_\tau = f^{-1}[[\tau]] \setminus U_\tau$. Notice that $f \upharpoonright G$ is continuous and since $X = G \cup \bigcup_\tau Z_\tau$ there is $\tau \in \omega^{<\omega}$ such that $Z_\tau \not\subseteq I_f$. Observe that $Z_\tau \subseteq \text{cl}(\bigcup_{\tau' \neq \tau, |\tau'|=|\tau|} f^{-1}[[\tau']])$ because if an open set $U \subseteq f^{-1}[[\tau]]$ is disjoint from $\bigcup_{\tau' \neq \tau, |\tau'|=|\tau|} f^{-1}[[\tau']]$ then $U \subseteq U_\tau$. Now put $C_\omega = [\tau]$ and pick any compact set with small diameter $X_\omega \subseteq Z_\tau$ such that $X_\omega \not\subseteq I_f$. $\square$

The construction. We begin with $X_\emptyset = X_\emptyset^0 = X$ and $C_\emptyset = \omega^\omega$. Without loss of generality assume that $X = X^*$. Suppose we have done $n - 1$ steps of the inductive construction up to $\tau \in (\omega + 1)^{<\omega}$. Let $|\tau| = l$ and $\sigma = \tau \upharpoonright (l - 1)$. There are three cases.

Case 1. The four points τ , $\pi(\tau)$, $r(\tau)$ and $r(\pi(\tau))$ are equal. So $\tau = (\omega, \dots, \omega)$ and $C_{\tau|n-1}$ and X_σ^{n-1} are already constructed. In this case we use Lemma 2 to find a clopen set C_τ and a compact set $X_\tau \subseteq X_\sigma^{n-1}$ of diameter $< |\tau|/3^{n+1}$ small enough so that no element of the anchors constructed so far is contained in X_τ . We put $X_\tau^n = X_\tau^*$, $X_\sigma^n = (X_\sigma^{n-1} \setminus f^{-1}[C_\tau])^*$ and $X_\rho^n = X_\rho^{n-1}$ for other $\rho < \tau$. By the assertion of Lemma 2 we still have $X_\tau^n \subseteq \text{cl}(X_\sigma^n)$. In this case we do not need to construct any new anchors.

Case 2. The two points $\pi(\tau)$ and $r(\tau)$ are equal but distinct from τ . Let $\delta = d(\tau, r(\tau))$. Since $X_{r(\tau)}^{n-1} \subseteq \text{cl}(X_\sigma^{n-1})$ by the inductive assumption, we may find finitely many sets $B_i \subseteq X_\sigma^{n-1}$, $i \leq k$ such that

- $h(\bigcup_i B'_i, X_{r(\tau)}) < \delta$ for any $B'_i \subseteq B_i$,
- $B_i \not\subseteq I_f$.

The second condition follows from $X_\sigma^{n-1} = (X_\sigma^{n-1})^*$. We may assume that for each clopen set $C \subseteq 2^\omega$ the set $B_i \cap f^{-1}[C]$ is either empty or outside of the ideal I_f .

We are going to find clopens $C_i \subseteq C_\sigma$, for $i \leq k$ such that $C_i \cap C_{r(\tau)} = \emptyset$ and then put $C_\tau = \bigcup_{i \leq k} C_i$, $X_\sigma^n = (X_\sigma^{n-1} \setminus \bigcup_i f^{-1}[C_i])^*$ and find $X_\tau \subseteq \bigcup_{i \leq k} B_i \cap f^{-1}[C_i]$. We will have to carefully define $X_{r(\tau)}^n$ so that $X_{r(\tau)}^n \subseteq \text{cl}(X_\sigma^n)$.

It is easy to see that for any $A \subseteq X_\sigma^{n-1}$

$$X_{r(\tau)}^{n-1} = X_{r(\tau)}^{n-1} \cap \text{cl}((X_\sigma^{n-1} \cap A)^*) \cup X_{r(\tau)}^{n-1} \cap \text{cl}((X_\sigma^{n-1} \cap A^c)^*)$$

so (putting $A = f^{-1}[C_\sigma \cap [(m, 0)]]$ for $m < \omega$) we may inductively on m pick binary sequences $\beta_i^m \in 2^m$, $i \leq k$ such that $f^{-1}[[\beta_i^m]] \cap B_i \neq \emptyset$ and

$$X_{r(\tau)}^{n-1} \cap \text{cl}((X_\sigma^{n-1} \setminus f^{-1}[\bigcup_{i \leq k} [\beta_i^m]])^*) \notin I_f.$$

We are going to carry on this construction up to some $m < \omega$ and put $X_\rho^n = (X_\rho^{n-1} \setminus \bigcup_{i \leq k} f^{-1}[[\beta_i^m]])^*$ for $\rho < \tau$, $\rho \not\geq r(\tau)$ and $X_\rho^n = (X_\rho^{n-1} \cap \text{cl}(X_\sigma^n))^*$ for $\rho < \tau$, $\rho \geq r(\tau)$. We must, however, take care that this does not destroy the existing anchors.

Since $f^{-1}[\{x\}] \in I_f$ for any $x \in 2^\omega$ and there are only finitely many elements of the existing anchors, we may pick $m < \omega$ and construct the sequences β_i^m so that for any element A of an anchor “below” $X_{r(\tau)}^{n-1}$ it is the case that $A \cap \text{cl}(f^{-1}[C_\sigma \setminus \bigcup_{i \leq k} [\beta_i^m]] \cap X_\sigma^{n-1}) \notin I_f$ and for any element A of other anchors $A \setminus (\bigcup_{i \leq k} f^{-1}[\beta_i^m]) \notin I_f$.

Once we have constructed the sequences β_i^m for $i \leq k$ we put $C_i = [\beta_i^m]$ and $C_\tau = \bigcup_i [\beta_i^m]$. Next we find I_f -positive compact sets X_i inside $B_i \cap f^{-1}[[\beta_i]]$, each of diameter $< 1/(3^{n+1}|\tau|)$.

If $\delta > \text{diam}(X_{\pi(\tau)}^{n-1})$ then we can pick one X_i as X_τ and then $h(X_\tau, X_{\pi(\tau)}^{n-1}) \leq 3h(X_\sigma^{n-1}, X_{\pi(\sigma)}^{n-1}) < 3^{|\tau|}\delta$. Otherwise, let $X_\tau = \bigcup_{i \leq k} X_i$ and then $\text{diam}(X_\tau) < 3\text{diam}(X_{\pi(\tau)}^{n-1}) \leq 3\text{diam}(X_{\pi(\tau)})$. Define $X_\tau^n = X_\tau^*$.

At this step we create anchors for the pair X_τ^n and $X_{r(\tau)}^n$ as well as for the tripples $X_\sigma^n, X_\rho^n, X_\tau^n$ for $\rho < \tau$.

Case 3. The two points $\pi(\tau), r(\tau)$ are distinct. Let $\delta = d(\tau, \pi(\tau))$. By coherence of the projections $\pi(\tau) \supseteq \pi(\sigma)$. By the inductive assumption we have $h(X_\sigma^{n-1}, X_{\pi(\sigma)}^{n-1}) < 3^{|\sigma|}\delta$. Using the existing anchor for the tripple $X_\sigma^{n-1}, X_{\pi(\sigma)}^{n-1}, X_{\pi(\tau)}^{n-1}$ let us find finitely many sets $B_i, i \leq k$ in X_σ such that

- $h(\bigcup_i B'_i, X_{\pi(\tau)}) < 3^{|\sigma|}\delta$ for any $B'_i \subseteq B_i$,
- $B_i \notin I_f$.

As before, we assume that for each clopen set $C \subseteq 2^\omega$ if $B_i \cap f^{-1}[C] \in I_f$ then it is empty. We have now two subcases, in analogy to the two previous cases.

Subcase 3.1. Suppose $\tau = r(\tau)$. Similarly as in Case 1, we use Lemma 2 to find $X_i \subseteq B_i$ and C_i for $i \leq k$. Put $C_\tau = \bigcup_{i \leq k} C_i$. If $\delta > \text{diam}(X_{\pi(\tau)}^{n-1})$ then we can pick one X_i as X_τ and then $h(X_\tau, X_{\pi(\tau)}^{n-1}) \leq 3h(X_\sigma^{n-1}, X_{\pi(\sigma)}^{n-1}) < 3^{|\tau|}\delta$. Otherwise, let $X_\tau = \bigcup_{i \leq k} X_i$ and then $\text{diam}(X_\tau) < 3\text{diam}(X_{\pi(\tau)}^{n-1}) \leq 3\text{diam}(X_{\pi(\tau)})$. Again, similarly as in Case 1, we put $X_\tau^n = (X_\tau^*)^*$, $X_\sigma^n = (X_\sigma^{n-1} \setminus f^{-1}[C_\tau])^*$, $X_\rho^n = X_\rho^{n-1}$ for other $\rho < \tau$.

Subcase 3.2. Suppose $\tau \neq r(\tau)$. Similarly as in Case 2, we find clopens C_i in ω^ω such that $X_{r(\tau)}^{n-1} \cap \text{cl}((f^{-1}[C_\sigma \setminus \bigcup_{i \leq k} C_i])^*) \notin I_f$ and no existing anchor is destroyed when we put $X_\rho^n = (X_\rho^{n-1} \setminus \bigcup_{i \leq k} f^{-1}[C_i])^*$ for $\rho < \tau, \rho \not\supseteq r(\tau)$ and $X_\rho^n = X_\rho^{n-1} \cap \text{cl}(X_\sigma^n)$ for $\rho < \tau, \rho \supseteq r(\tau)$.

Next we find I_f -positive compact sets $X_i \subseteq B_i \cap f^{-1}[C_i]$ each of diameter $< 1/(3^{|\tau|+1}|\tau|)$. As previously, if $\delta > \text{diam}(X_{\pi(\tau)}^{n-1})$ then we can pick one X_i as X_τ and then $h(X_\tau, X_{\pi(\tau)}^{n-1}) \leq 3h(X_\sigma^{n-1}, X_{\pi(\sigma)}^{n-1}) < 3^{|\tau|}\delta$. Otherwise, let $X_\tau = \bigcup_{i \leq k} X_i$ and then $\text{diam}(X_\tau) < 3\text{diam}(X_{\pi(\tau)}^{n-1}) \leq 3\text{diam}(X_{\pi(\tau)})$. Again, we put $X_\tau^n = (X_\tau^*)^*$,

In Case 3 we construct the same anchors as in Case 2.

This ends the construction and the entire proof. $\square$

Theorem 5. *If $f : X \rightarrow \omega^\omega$ is not σ -continuous then there exist topological embeddings φ and ψ such that the following diagram commutes:*

$$\begin{array}{ccc} \omega^\omega & \xrightarrow{\psi} & \omega^\omega \\ \uparrow P & & \uparrow f \\ (\omega + 1)^\omega & \xrightarrow{\varphi} & X \end{array}$$

Proof. By Theorem 4 we have ψ and φ such that ψ is 1-1 open. But as a Borel function it is continuous on a dense G_δ set $G \subseteq \omega^\omega$. On the other hand by the properties of the function P $X \in I_P$ implies $P[X]$ is meager. So $P^{-1}[G] \notin I_P$ and the problem reduces to the restriction of the function P . This, however, has been proved in [2] (Corollary 2). So we get the following diagram:

$$\begin{array}{ccccc} \omega^\omega & \xrightarrow{\psi'} & G & \xrightarrow{\psi} & \omega^\omega \\ \uparrow P & & \uparrow P|_G & & \uparrow f \\ (\omega + 1)^\omega & \xrightarrow{\varphi'} & P^{-1}[G] & \xrightarrow{\varphi} & X \end{array}$$

which ends the proof. $\square$

REFERENCES

- [1] Cichoń J., Morayne M., Pawlikowski J. and Solecki S., *Decomposing Baire functions*, Journal of Symbolic Logic, Vol. 56, Issue 4, 1991, pp. 1273-1283
- [2] Sabok M., *σ -continuity and related forcings*, submitted,
- [3] Solecki S., *Decomposing Borel sets and functions and the structure of Baire class 1 functions*, Journal of the American Mathematical Society, Vol. 11, No. 3, 1998, pp. 521-550
- [4] Zapletal J., *Descriptive Set Theory and Definable Forcing*, Memoirs of the American Mathematical Society, 2004
- [5] Zapletal J., *Forcing Idealized*, Cambridge Tracts in Mathematics 174, 2008

MATHEMATICAL INSTITUTE, WROCLAW UNIVERSITY, PL. GRUNWALDZKI 2/4,
50-384 WROCLAW, POLAND

E-mail address: sabok@math.uni.wroc.pl